සියලුම හිමිකම් ඇවිරිණි/மුඟුப் பதிப்புரிமையுடையது/ $All\ Rights\ Reserved$ ]

ලි ලංකා විභාග දෙපාර්තමේන්තුව ලි ලංකා විභාග දෙපාර්තමේන්තුව දි ලංකා විභාග දෙපාර්තමේන්තුව ලි ලංකා විභාග දෙපාර්තමේන්තුව இலங்கைப் பரீட்சைத் திணைக்களம் இலங்கைப் பரீட்சைத் திணைக்களம் இலங்கைப் பரீட்சைத் திணைக்களம் Department of Examinations, Sri Lanka Department o**இலங்கைப் இலங்கைப் பரீட்**சைத் நினைக்களம் ලි ලංකා විභාග දෙපාර්තමේන්තුව ලි ලංකා විභාග දෙපාර්තමේන්තුව ලී ලංකා විභාග දෙපාර්තමේන්තුව

අධායන පොදු සහතික පතු (උසස් පෙළ) විභාගය, 2024 கல்விப் பொதுத் தராதரப் பத்திர (உயர் தர)ப் பரீட்சை, 2024 General Certificate of Education (Adv. Level) Examination, 2024

උසස් ගණිතය II உயர் கணிதம் II Higher Mathematics II



පැය තුනයි மூன்று மணித்தியாலம் Three hours

Use additional reading time to go through the question paper, select the questions you will answer and decide which of them you will prioritise.

$\overline{}$						
Index	Number					
			100	0.7	1.8 % []	

#### Instructions:

- \* This question paper consists of two parts.

  Part A (Questions 1 10) and Part B (Questions 11 17).
- \* Part A:

  Answer all questions. Write your answers to each question in the space provided. You may use additional sheets if more space is needed.
- \* Part B:

  Answer five questions only. Write your answers on the sheets provided.
- \* At the end of the time allotted, tie the answer scripts of the two parts together so that Part A is on top of Part B and hand them over to the supervisor.
- \* You are permitted to remove only Part B of the question paper from the Examination Hall.
- \* Statistical Tables will be provided.
- \* g denotes the acceleration due to gravity.

### For Examiners' Use only

Part	Question No.	Marks			
	1		4		
	2				
	3				
	4				
A	5				
	6				
	7				
	8				
	9				
	10				
	11				
	12				
ъ	13		-		
В	14	* .	_		
	15				
	16				
	17				
	Total				

	Total	
In Numbers	<u> </u>	
In Words		

Marking Examiner

Checked by:

2

Supervised by:

# Part A

	Let $\mathbf{F} = \mathbf{i} + 2\mathbf{j} + \alpha\mathbf{k}$ , where $\alpha \in \mathbf{r} = a\mathbf{i} + \mathbf{j} + \lambda(\mathbf{i} + 2\mathbf{j} + \alpha\mathbf{k})$ and the where $a, b \in \mathbb{R}$ . If the vector materials	he force -	$-\mathbf{F}$ acting a	along the li	ine $r = i + b$	$b\mathbf{j} + \mathbf{k} + \mu$	u(1 + 2j +	$\alpha \mathbf{K}$ ),
							• • • • • • • • • • • • • • • • • • • •	
			•••••		*****			
						• • • • • • • • • • • • • • • • • • • •		
	· · · · · · · · · · · · · · · · · · ·	•••••	 	•••••				
				************			**************************************	•••
								••••
-					****			••••
				·				•••
		gregarians server	4 6 6 3 4 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6					**************************************
			**********					
								••••
								· · · ·
								····
		and the second		20 P. 15				
		(				******		• • • •
2	Relative to a fived origin O	the positi	ion vectors	of three i	ooints A. B	and $C$	are 2i+i	 j + k.
2.	Relative to a fixed origin $O$ , is $2\mathbf{i} + \mathbf{k}$ and $2\mathbf{i} - \mathbf{i} - \mathbf{k}$ , respect	the positi	ion vectors $\overrightarrow{AR} \times \overrightarrow{AI}$	of three t	points A, B	and C	are 2i + j	 <b>j + k,</b> cular
2.	Relative to a fixed origin $O$ , $i-2j+k$ and $2i-j-k$ , respect to the plane containing $A$ , $B$ a	tively. Fin	on vectors and $\overrightarrow{AB} \times \overrightarrow{AC}$	of three $\vec{C}$ and hence	ooints A, B	and C	are 2 <b>i</b> + <b>j</b> perpendi	 <b>j + k,</b> cular
2.	i-2j+k and $2i-j-k$ , respect	tively. Fin	ion vectors at $\overrightarrow{AB} \times \overrightarrow{A0}$	of three $\vec{z}$ and hence	ooints A, B	and C	are 2i + j perpendi	j + k, cular
2.	i-2j+k and $2i-j-k$ , respect	tively. Fin	ion vectors id $\overrightarrow{AB} \times \overrightarrow{AC}$	of three properties and hence	ooints A, B	and C	are 2i + j perpendi	 j + k, cular 
2.	i-2j+k and $2i-j-k$ , respect	tively. Fin	ion vectors at $\overrightarrow{AB} \times \overrightarrow{AC}$	of three $\vec{C}$ and hence	ooints A, B	and C	are 2i + j perpendi	j + k, cular 
2.	i-2j+k and $2i-j-k$ , respect	tively. Fin	on vectors at $\overrightarrow{AB} \times \overrightarrow{AC}$	of three t	oints A, B	and C	are 2i + j perpendi	j + k, cular 
2.	i-2j+k and $2i-j-k$ , respect	tively. Fin	on vectors at $\overrightarrow{AB} \times \overrightarrow{A0}$	of three t	oints A, B	and C	are 2i + j	j + <b>k</b> , cular
2.	i-2j+k and $2i-j-k$ , respect	tively. Fin	on vectors at $\overrightarrow{AB} \times \overrightarrow{A0}$	of three t	oints A, B	and C	are 2i + j	j + k, cular
2.	i-2j+k and $2i-j-k$ , respect	tively. Fin	on vectors at $\overrightarrow{AB} \times \overrightarrow{A0}$	of three t	ooints A, B	and C	are 2i + j	j + k, cular 
2.	i-2j+k and $2i-j-k$ , respect	tively. Fin	on vectors of $\overrightarrow{AB} \times \overrightarrow{AC}$	of three t	ooints A, B	and C	are 2i + j perpendi	j + k, cular
2.	i-2j+k and $2i-j-k$ , respect	tively. Fin	on vectors of $\overrightarrow{AB} \times \overrightarrow{AC}$	of three t	oints A, B	and C	are 2i + j perpendi	j + k, cular
2.	i-2j+k and $2i-j-k$ , respect	tively. Fin	on vectors $d \overline{AB} \times \overline{AC}$	of three t	oints A, B	and C	are 2i + j perpendi	j + k, cular
2.	i-2j+k and $2i-j-k$ , respect	tively. Fin	on vectors $d \overline{AB} \times \overline{AC}$	of three t	oints A, B	and C it vector	are 2i + j perpendi	j+k, cular
2.	i-2j+k and $2i-j-k$ , respect	tively. Fin	on vectors $d \overline{AB} \times \overline{AC}$	of three t	oints A, B	and C it vector	are 2i + j perpendi	j + k, cular
2.	i-2j+k and $2i-j-k$ , respect	tively. Fin	on vectors $d \overline{AB} \times \overline{AC}$	of three the and hence	oints A, B	and C it vector	are 2i + j perpendi	j + k, cular

[see page four

	***************************************	· · · · · · · · · · · · · · · · · · ·
······		• • • • • • • • • • • • • • • • • • • •
••••		
	anne ann an ann ann ann an an an an an an an	

The velocity vector of a particle P at time t is given by  $\mathbf{v}(t) = (\sin t + \cos t)\mathbf{i} + (t + t^2)\mathbf{j} + e^t\mathbf{k}$ . It is also given that at t = 0, the position vector of P is equal to the acceleration vector of P. Find the position vector of P at time t.

J.	A smooth sphere $P$ of mass $m$ , moving at speed $u$ , collides with an identical sphere $Q$ which is initially at rest. Immediately before the collision, the velocity of $P$ makes an angle $\theta$ with the line of centres. Immediately after the collision, the velocity of $P$ makes an angle $\phi$ with the line of centres (see the figure). The coefficient of restitution $2 \tan \theta$	· <b></b>
	between the spheres is $e(e < 1)$ . Show that $\tan \phi = \frac{2 \tan \theta}{1 - e}$ .	
		••
		••
. in		•
7.		••
		••
All A		•••
		•••
6	prings (2011年) - Angle (1911年)	
<b>U.</b>	A uniform solid right circular cylinder of mass $M$ has radius $4a$ and height $6a$ . It is given that the moment of inertia of the cylinder	7
0.	and height 6a. It is given that the moment of inertia of the cylinder about an axis through the centre of mass perpendicular to the axis of	
0.	and height $6a$ . It is given that the moment of inertia of the cylinder about an axis through the centre of mass perpendicular to the axis of the cylinder is $7Ma^2$ . The cylinder can rotate about a fixed smooth horizontal axis through a point $A$ on the circumference of one end of	
	and height 6a. It is given that the moment of inertia of the cylinder about an axis through the centre of mass perpendicular to the axis of the cylinder is $7Ma^2$ . The cylinder can rotate about a fixed smooth horizontal axis through a point A on the circumference of one end of the cylinder. The axis of rotation is tangential to this circumference. The cylinder is held at rest with axis of the cylinder horizontal and	
	and height $6a$ . It is given that the moment of inertia of the cylinder about an axis through the centre of mass perpendicular to the axis of the cylinder is $7Ma^2$ . The cylinder can rotate about a fixed smooth horizontal axis through a point $A$ on the circumference of one end of the cylinder. The axis of rotation is tangential to this circumference. The cylinder is held at rest with axis of the cylinder horizontal and at a height $4a$ above the point $A$ (see the figure). Show that, if the	
受ける できます かんしゅう かんしゅう	and height $6a$ . It is given that the moment of inertia of the cylinder about an axis through the centre of mass perpendicular to the axis of the cylinder is $7Ma^2$ . The cylinder can rotate about a fixed smooth horizontal axis through a point $A$ on the circumference of one end of the cylinder. The axis of rotation is tangential to this circumference. The cylinder is held at rest with axis of the cylinder horizontal and at a height $4a$ above the point $A$ (see the figure). Show that, if the cylinder is released, the angular speed at the instant when the axis of	
気と感染を含むした。 これのであればない	and height $6a$ . It is given that the moment of inertia of the cylinder about an axis through the centre of mass perpendicular to the axis of the cylinder is $7Ma^2$ . The cylinder can rotate about a fixed smooth horizontal axis through a point $A$ on the circumference of one end of the cylinder. The axis of rotation is tangential to this circumference. The cylinder is held at rest with axis of the cylinder horizontal and at a height $4a$ above the point $A$ (see the figure). Show that, if the	
のであるできます。 こうこうから はずべつ	and height $6a$ . It is given that the moment of inertia of the cylinder about an axis through the centre of mass perpendicular to the axis of the cylinder is $7Ma^2$ . The cylinder can rotate about a fixed smooth horizontal axis through a point $A$ on the circumference of one end of the cylinder. The axis of rotation is tangential to this circumference. The cylinder is held at rest with axis of the cylinder horizontal and at a height $4a$ above the point $A$ (see the figure). Show that, if the cylinder is released, the angular speed at the instant when the axis of	
教と考えている。 ・	and height $6a$ . It is given that the moment of inertia of the cylinder about an axis through the centre of mass perpendicular to the axis of the cylinder is $7Ma^2$ . The cylinder can rotate about a fixed smooth horizontal axis through a point $A$ on the circumference of one end of the cylinder. The axis of rotation is tangential to this circumference. The cylinder is held at rest with axis of the cylinder horizontal and at a height $4a$ above the point $A$ (see the figure). Show that, if the cylinder is released, the angular speed at the instant when the axis of	
がなった。 できない かんしゅう かんしゅ かんしゅう かんしゅう かんしゅう かんしゅう かんしゅう かんしゃ かんしゅん かんしゃ かんしゃ かんしゃ かんしゃ かんしゃ かんしゃ かんしゃ かんし	and height $6a$ . It is given that the moment of inertia of the cylinder about an axis through the centre of mass perpendicular to the axis of the cylinder is $7Ma^2$ . The cylinder can rotate about a fixed smooth horizontal axis through a point $A$ on the circumference of one end of the cylinder. The axis of rotation is tangential to this circumference. The cylinder is held at rest with axis of the cylinder horizontal and at a height $4a$ above the point $A$ (see the figure). Show that, if the cylinder is released, the angular speed at the instant when the axis of the cylinder is vertical is $\sqrt{\frac{7g}{16a}}$ .	
がある。 こうかん こうかん はっぱん しゅうしゅう こうかん はっぱん こうかん はっぱん こうかん こうかん こうかん こうかん こうかん こうかん こうかん こうか	and height $6a$ . It is given that the moment of inertia of the cylinder about an axis through the centre of mass perpendicular to the axis of the cylinder is $7Ma^2$ . The cylinder can rotate about a fixed smooth horizontal axis through a point $A$ on the circumference of one end of the cylinder. The axis of rotation is tangential to this circumference. The cylinder is held at rest with axis of the cylinder horizontal and at a height $4a$ above the point $A$ (see the figure). Show that, if the cylinder is released, the angular speed at the instant when the axis of the cylinder is vertical is $\sqrt{\frac{7g}{16a}}$ .	
	and height $6a$ . It is given that the moment of inertia of the cylinder about an axis through the centre of mass perpendicular to the axis of the cylinder is $7Ma^2$ . The cylinder can rotate about a fixed smooth horizontal axis through a point $A$ on the circumference of one end of the cylinder. The axis of rotation is tangential to this circumference. The cylinder is held at rest with axis of the cylinder horizontal and at a height $4a$ above the point $A$ (see the figure). Show that, if the cylinder is released, the angular speed at the instant when the axis of the cylinder is vertical is $\sqrt{\frac{7g}{16a}}$ .	
大型のできます。 のでは、	and height $6a$ . It is given that the moment of inertia of the cylinder about an axis through the centre of mass perpendicular to the axis of the cylinder is $7Ma^2$ . The cylinder can rotate about a fixed smooth horizontal axis through a point $A$ on the circumference of one end of the cylinder. The axis of rotation is tangential to this circumference. The cylinder is held at rest with axis of the cylinder horizontal and at a height $4a$ above the point $A$ (see the figure). Show that, if the cylinder is released, the angular speed at the instant when the axis of the cylinder is vertical is $\sqrt{\frac{7g}{16a}}$ .	
大学 Andrew Andr	and height $6a$ . It is given that the moment of inertia of the cylinder about an axis through the centre of mass perpendicular to the axis of the cylinder is $7Ma^2$ . The cylinder can rotate about a fixed smooth horizontal axis through a point $A$ on the circumference of one end of the cylinder. The axis of rotation is tangential to this circumference. The cylinder is held at rest with axis of the cylinder horizontal and at a height $4a$ above the point $A$ (see the figure). Show that, if the cylinder is released, the angular speed at the instant when the axis of the cylinder is vertical is $\sqrt{\frac{7g}{16a}}$ .	
大学のできます。 The Control of the Contro	and height $6a$ . It is given that the moment of inertia of the cylinder about an axis through the centre of mass perpendicular to the axis of the cylinder is $7Ma^2$ . The cylinder can rotate about a fixed smooth horizontal axis through a point $A$ on the circumference of one end of the cylinder. The axis of rotation is tangential to this circumference. The cylinder is held at rest with axis of the cylinder horizontal and at a height $4a$ above the point $A$ (see the figure). Show that, if the cylinder is released, the angular speed at the instant when the axis of the cylinder is vertical is $\sqrt{\frac{7g}{16a}}$ .	
大学のできます。 The Control of Manager Control of Manage	and height $6a$ . It is given that the moment of inertia of the cylinder about an axis through the centre of mass perpendicular to the axis of the cylinder is $7Ma^2$ . The cylinder can rotate about a fixed smooth horizontal axis through a point $A$ on the circumference of one end of the cylinder. The axis of rotation is tangential to this circumference. The cylinder is held at rest with axis of the cylinder horizontal and at a height $4a$ above the point $A$ (see the figure). Show that, if the cylinder is released, the angular speed at the instant when the axis of the cylinder is vertical is $\sqrt{\frac{7g}{16a}}$ .	
	and height $6a$ . It is given that the moment of inertia of the cylinder about an axis through the centre of mass perpendicular to the axis of the cylinder is $7Ma^2$ . The cylinder can rotate about a fixed smooth horizontal axis through a point $A$ on the circumference of one end of the cylinder. The axis of rotation is tangential to this circumference. The cylinder is held at rest with axis of the cylinder horizontal and at a height $4a$ above the point $A$ (see the figure). Show that, if the cylinder is released, the angular speed at the instant when the axis of the cylinder is vertical is $\sqrt{\frac{7g}{16a}}$ .	
のでは、「Andrews Andrews	and height $6a$ . It is given that the moment of inertia of the cylinder about an axis through the centre of mass perpendicular to the axis of the cylinder is $7Ma^2$ . The cylinder can rotate about a fixed smooth horizontal axis through a point $A$ on the circumference of one end of the cylinder. The axis of rotation is tangential to this circumference. The cylinder is held at rest with axis of the cylinder horizontal and at a height $4a$ above the point $A$ (see the figure). Show that, if the cylinder is released, the angular speed at the instant when the axis of the cylinder is vertical is $\sqrt{\frac{7g}{16a}}$ .	

	A student randomly guesses answers to a choices, of which only one choice is con independent of his choice for any other	rect. Assume that the	student's choice for a	question is
	Calculate the probability that	<b>\$</b>		
	(i) answers to the first three questions a	re all correct.		
	(ii) at most two of the answers to the fi		e correct.	e.
	(ii) at most two of the planta to the	1		
				*
	\$			
	·		Part Control of the C	
	•••••			
			Andrews	
	. *************************************			
		A Section 1		
2		Albaria Barrana and Andr		
	The second of th			وبير وسيد
				· • • • • • • • • • • • • • • • • • • •
	• • • • • • • • • • • • • • • • • • •			
8.	The lifetime of a certain machine follow machine does not fail during the first y machine will operate without failure for Given that the machine has not failed du it will continue to operate without failure	year of operation is of more than two years.  uring its first year of o	0.7. Find the probabilit	y that the
		The second secon		
•				
•				
-				
•				

			. H U > *	١.					
	$F(y) = \int$		$   \text{if } 0 \le x \le 3 \\   \text{if } x > 3 $						
	$I^{r}(\lambda)^{-}$	Δ ,	, if $x>3$ , otherwise.						
	-				C V				
Find th	ie value c	of $K$ , the	median of $X$	, and the	mean of $X$ .				
				<sub>.</sub>		*********			
						9			
• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •			,					
	<i>,</i>	• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •				·,		
					. '. 				
					· · · · · · · · · · · · · · · · · · ·				
• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •				••••••		**********	••••
			· · · · · · · · · · · · · · · · · · ·	• • • • • • • • • • • • • • • • • • • •					
								*	
	•••••			· · · · · · · · · · · · · · · · · · ·				R & G = -	
				11.5. 12부 (취임 2개합.편		: 			
						• • • • • • • • • • • • • • • • • • • •			••••
•••••		enteriores de la composición de la com La composición de la		~ A 4 1			• • • • • • • • • • • • • • • • • • • •		• • • • • •
		· <u>.</u>		· ·	t was a second		,		
		* * * * * * * * * * * * * * * * * * *			Kan	•••••			-
		1444							OB A
The pro	obability	mass func	ction of a di	screte ran		e $X$ is as			•••••
The pro		<del></del>	11 11 11 11 11 11 11	<del> </del>	dom variabl	e X is as			
	x	0	1	2	dom variabl	e <i>X</i> is as			
P	X $P(X=x)$	0 0.1	0.2	2 k	dom variabl	e <i>X</i> is as 4 0.1			
Pind th	$\begin{array}{c c} x & \\ P(X=x) & \\ \text{ne value of } \end{array}$	$\begin{array}{c} 0 \\ \hline 0.1 \\ \text{of } k, \text{ the } \end{array}$	$\begin{array}{c c} & 1 \\ \hline & 0.2 \\ \hline \end{array}$ mean of $X$ a	$\frac{2}{k}$ and the variation	$\begin{array}{ c c c } \hline \text{dom variabl} \\ \hline & 3 \\ \hline & 0.2 \\ \hline \text{ariance of } X \\ \hline \end{array}$	$ \begin{array}{c}                                     $	follows:		
Pind th	$\begin{array}{c c} x & \\ P(X=x) & \\ \text{ne value of } \end{array}$	$\begin{array}{c} 0 \\ \hline 0.1 \\ \text{of } k, \text{ the } \end{array}$	0.2	$\frac{2}{k}$ and the variation	$\begin{array}{ c c c } \hline \text{dom variabl} \\ \hline & 3 \\ \hline & 0.2 \\ \hline \text{ariance of } X \\ \hline \end{array}$	$ \begin{array}{c}                                     $	follows:	a.	
Pind th	$\begin{array}{c c} x & \\ P(X=x) & \\ \text{ne value of } \end{array}$	$\begin{array}{c} 0 \\ \hline 0.1 \\ \text{of } k, \text{ the } \end{array}$	$\begin{array}{c c} & 1 \\ \hline & 0.2 \\ \hline \end{array}$ mean of $X$ a	$\frac{2}{k}$ and the variation	$\begin{array}{ c c c } \hline \text{dom variabl} \\ \hline & 3 \\ \hline & 0.2 \\ \hline \text{ariance of } X \\ \hline \end{array}$	$ \begin{array}{c}                                     $	follows:	a.	
Pind th	$\begin{array}{c c} x & \\ P(X=x) & \\ \text{ne value of } \end{array}$	$\begin{array}{c} 0 \\ \hline 0.1 \\ \text{of } k, \text{ the } \end{array}$	$\begin{array}{c c} & 1 \\ \hline & 0.2 \\ \hline \end{array}$ mean of $X$ a	$\frac{2}{k}$ and the variation	$\begin{array}{ c c c } \hline \text{dom variabl} \\ \hline & 3 \\ \hline & 0.2 \\ \hline \text{ariance of } X \\ \hline \end{array}$	$ \begin{array}{c}                                     $	follows:	a.	
Pind th	$\begin{array}{c c} x & \\ P(X=x) & \\ \text{ne value of } \end{array}$	$\begin{array}{c} 0 \\ \hline 0.1 \\ \text{of } k, \text{ the } \end{array}$	$\begin{array}{c c} & 1 \\ \hline & 0.2 \\ \hline \end{array}$ mean of $X$ a	$\frac{2}{k}$ and the variation	$\begin{array}{ c c c } \hline \text{dom variabl} \\ \hline & 3 \\ \hline & 0.2 \\ \hline \text{ariance of } X \\ \hline \end{array}$	$ \begin{array}{c}                                     $	follows:	a.	
Pind th	$\begin{array}{c c} x & \\ P(X=x) & \\ \text{ne value of } \end{array}$	$\begin{array}{c} 0 \\ \hline 0.1 \\ \text{of } k, \text{ the } \end{array}$	$\begin{array}{c c} & 1 \\ \hline & 0.2 \\ \hline \end{array}$ mean of $X$ a	$\frac{2}{k}$ and the variation	$\begin{array}{ c c c } \hline \text{dom variabl} \\ \hline & 3 \\ \hline & 0.2 \\ \hline \text{ariance of } X \\ \hline \end{array}$	$ \begin{array}{c}                                     $	follows:	a.	
Pind th	$\begin{array}{c c} x & \\ P(X=x) & \\ \text{ne value of } \end{array}$	$\begin{array}{c} 0 \\ \hline 0.1 \\ \text{of } k, \text{ the } \end{array}$	$\begin{array}{c c} & 1 \\ \hline & 0.2 \\ \hline \end{array}$ mean of $X$ a	$\frac{2}{k}$ and the variation	$\begin{array}{ c c c } \hline \text{dom variabl} \\ \hline & 3 \\ \hline & 0.2 \\ \hline \text{ariance of } X \\ \hline \end{array}$	$ \begin{array}{c}                                     $	follows:	a.	
Pind th	$\begin{array}{c c} x & \\ P(X=x) & \\ \text{ne value of } \end{array}$	$\begin{array}{c} 0 \\ \hline 0.1 \\ \text{of } k, \text{ the } \end{array}$	$\begin{array}{c c} & 1 \\ \hline & 0.2 \\ \hline \end{array}$ mean of $X$ a	$\frac{2}{k}$ and the variation	$\begin{array}{ c c c } \hline \text{dom variabl} \\ \hline & 3 \\ \hline & 0.2 \\ \hline \text{ariance of } X \\ \hline \end{array}$	$ \begin{array}{c}                                     $	follows:	a.	
Pind th	$\begin{array}{c c} x & \\ P(X=x) & \\ \text{ne value of } \end{array}$	$\begin{array}{c} 0 \\ \hline 0.1 \\ \text{of } k, \text{ the } \end{array}$	$\begin{array}{c c} & 1 \\ \hline & 0.2 \\ \hline \end{array}$ mean of $X$ a	$\frac{2}{k}$ and the variation	$\begin{array}{ c c c } \hline \text{dom variabl} \\ \hline & 3 \\ \hline & 0.2 \\ \hline \text{ariance of } X \\ \hline \end{array}$	$ \begin{array}{c}                                     $	follows:	a.	
Pind th	$\begin{array}{c c} x & \\ P(X=x) & \\ \text{ne value of } \end{array}$	$\begin{array}{c} 0 \\ \hline 0.1 \\ \text{of } k, \text{ the } \end{array}$	$\begin{array}{c c} & 1 \\ \hline & 0.2 \\ \hline \end{array}$ mean of $X$ a	$\frac{2}{k}$ and the variation	$\begin{array}{ c c c } \hline \text{dom variabl} \\ \hline & 3 \\ \hline & 0.2 \\ \hline \text{ariance of } X \\ \hline \end{array}$	$ \begin{array}{c}                                     $	follows:	a.	
Pind th	$\begin{array}{c c} x & \\ P(X=x) & \\ \text{ne value of } \end{array}$	$\begin{array}{c} 0 \\ \hline 0.1 \\ \text{of } k, \text{ the } \end{array}$	$\begin{array}{c c} & 1 \\ \hline & 0.2 \\ \hline \end{array}$ mean of $X$ a	$\frac{2}{k}$ and the variation	$\begin{array}{ c c c } \hline \text{dom variabl} \\ \hline & 3 \\ \hline & 0.2 \\ \hline \text{ariance of } X \\ \hline \end{array}$	$ \begin{array}{c}                                     $	follows:	a.	
Pind th	$\begin{array}{c c} x & \\ P(X=x) & \\ \text{ne value of } \end{array}$	$\begin{array}{c} 0 \\ \hline 0.1 \\ \text{of } k, \text{ the } \end{array}$	$\begin{array}{c c} & 1 \\ \hline & 0.2 \\ \hline \end{array}$ mean of $X$ a	$\frac{2}{k}$ and the variation	$\begin{array}{ c c c } \hline \text{dom variabl} \\ \hline & 3 \\ \hline & 0.2 \\ \hline \text{ariance of } X \\ \hline \end{array}$	$ \begin{array}{c}                                     $	follows:	a.	
Pind th	$\begin{array}{c c} x & \\ P(X=x) & \\ \text{ne value of } \end{array}$	$\begin{array}{c} 0 \\ \hline 0.1 \\ \text{of } k, \text{ the } \end{array}$	$\begin{array}{c c} & 1 \\ \hline & 0.2 \\ \hline \end{array}$ mean of $X$ a	$\frac{2}{k}$ and the variation	$\begin{array}{ c c c } \hline \text{dom variabl} \\ \hline & 3 \\ \hline & 0.2 \\ \hline \text{ariance of } X \\ \hline \end{array}$	$ \begin{array}{c}                                     $	follows:	a.	
Pind th	$\begin{array}{c c} x & \\ P(X=x) & \\ \text{ne value of } \end{array}$	$\begin{array}{c} 0 \\ \hline 0.1 \\ \text{of } k, \text{ the } \end{array}$	$\begin{array}{c c} & 1 \\ \hline & 0.2 \\ \hline \end{array}$ mean of $X$ a	$\frac{2}{k}$ and the variation	$\begin{array}{ c c c } \hline \text{dom variabl} \\ \hline & 3 \\ \hline & 0.2 \\ \hline \text{ariance of } X \\ \hline \end{array}$	$ \begin{array}{c}                                     $	follows:	a.	
Pind th	$\begin{array}{c c} x & \\ P(X=x) & \\ \text{ne value of } \end{array}$	$\begin{array}{c} 0 \\ \hline 0.1 \\ \text{of } k, \text{ the } \end{array}$	$\begin{array}{c c} & 1 \\ \hline & 0.2 \\ \hline \end{array}$ mean of $X$ a	$\frac{2}{k}$ and the variation	$\begin{array}{ c c c } \hline \text{dom variabl} \\ \hline & 3 \\ \hline & 0.2 \\ \hline \text{ariance of } X \\ \hline \end{array}$	$ \begin{array}{c}                                     $	follows:	a.	
Pind th	$\begin{array}{c c} x & \\ P(X=x) & \\ \text{ne value of } \end{array}$	$\begin{array}{c} 0 \\ \hline 0.1 \\ \text{of } k, \text{ the } \end{array}$	$\begin{array}{c c} & 1 \\ \hline & 0.2 \\ \hline \end{array}$ mean of $X$ a	$\frac{2}{k}$ and the variation	$\begin{array}{ c c c } \hline \text{dom variabl} \\ \hline & 3 \\ \hline & 0.2 \\ \hline \text{ariance of } X \\ \hline \end{array}$	$ \begin{array}{c}                                     $	follows:	a.	
P Find the	$\begin{array}{c c} x & \\ P(X=x) & \\ \text{ne value of } \end{array}$	$\begin{array}{c} 0 \\ \hline 0.1 \\ \text{of } k, \text{ the } \end{array}$	$\begin{array}{c c} & 1 \\ \hline & 0.2 \\ \hline \end{array}$ mean of $X$ a	$\frac{2}{k}$ and the variation	$\begin{array}{ c c c } \hline \text{dom variabl} \\ \hline & 3 \\ \hline & 0.2 \\ \hline \text{ariance of } X \\ \hline \end{array}$	$ \begin{array}{c}                                     $	follows:	a.	
P Find the	$\begin{array}{c c} x & \\ P(X=x) & \\ \text{ne value of } \end{array}$	$\begin{array}{c} 0 \\ \hline 0.1 \\ \text{of } k, \text{ the } \end{array}$	$\begin{array}{c c} & 1 \\ \hline & 0.2 \\ \hline \end{array}$ mean of $X$ a	$\frac{2}{k}$ and the variation	$\begin{array}{ c c c } \hline \text{dom variabl} \\ \hline & 3 \\ \hline & 0.2 \\ \hline \text{ariance of } X \\ \hline \end{array}$	$ \begin{array}{c}                                     $	follows:	a.	
Pind th	$\begin{array}{c c} x & \\ P(X=x) & \\ \text{ne value of } \end{array}$	$\begin{array}{c} 0 \\ \hline 0.1 \\ \text{of } k, \text{ the } \end{array}$	$\begin{array}{c c} & 1 \\ \hline & 0.2 \\ \hline \end{array}$ mean of $X$ a	$\frac{2}{k}$ and the variation	$\begin{array}{ c c c } \hline \text{dom variabl} \\ \hline & 3 \\ \hline & 0.2 \\ \hline \text{ariance of } X \\ \hline \end{array}$	$ \begin{array}{c}                                     $	follows:	a.	

ដ៏ជទ្ធ២ សិចិត្តថិ កុរខិបិសី /(ហ្គ $\psi$ ប់ ប្រង់ប្បព្រិស្តាយប្រាក្សា/ $All\ Rights\ Reserved$ 

ලි ලංකා විහාශ දෙපාර්තමේන්තුව ලි ලංකා විභාග දෙපාර්තමේක ලියකුලෙනුණුකු දෙප්තුර්තමේන්තුව විභාග දෙපාර්තමේන්තුව ලි ලංකා විභාග දෙපාර්තමේන්තුව මූහත්කෙන්ට பර්ධකපති නිකාක්ෂයගැට මුහත්කෙන්ට පරිධක් නිකාක්ෂයගැට මුහත්කෙන්ට පරිධක් නිකාක්ෂයගැට මුහත්කෙන්ට පරිධක් නි Department of Examinations, Sri Lanka Department o **මු බැඩාකාස්ප් S/ පාර්තමේ**න්තුව මුහත්කෙන්ට පරිධක්වේ මුහත්කෙන්ට පරිධක්වේ මුහත්කෙන්ට පරිධක්වේ පරිධක්වේ මුහත්කෙන්ට පරිධක්වේ සිට පරිධක්වේ මුහත්කෙන්ට පරිධක්වේ මුහත්කෙන්ට පරිධක්වේ සිට පරිධක්වේ පරිධක්වේ සිට ප

අධායන පොදු සහතික පතු (උසස් පෙළ) විභාගය, 2024 සහ්ඛා් பொதுத் தராதரப் பத்திர (உயர் தர)ப் பரீட்சை, 2024 General Certificate of Education (Adv. Level) Examination, 2024

උසස් ගණිතය II உயர் கணிதம் II Higher Mathematics II



### Part B

- \* Answer five questions only.
- 11. (a) A force **F** produces a moment of  $2\mathbf{i} 2\mathbf{j} + 3\mathbf{k}$  about the origin O. The line of action of **F** lies on a plane perpendicular to the x axis and passes through the point  $A \equiv (1, 1, 0)$ .
  - (i) Determine F.
  - (ii) It is given that the force **F** has zero moment about the point  $B \equiv (p, 0, q)$ . Find the values of p and q.
  - (b) Relative to a fixed origin O, points A and B of a rigid body have position vectors  $b\mathbf{i}$  and  $3\mathbf{i} + 4\mathbf{j} + 12\mathbf{k}$  respectively. A force  $\mathbf{F}$ , acting on the body, passes through O and its line of action is parallel to AB. Forces  $b\mathbf{i} 8\mathbf{j} + 8\mathbf{k}$  and  $p\mathbf{k}$ , also acting on the body, pass through points with position vectors  $b\mathbf{f}$  and  $3\mathbf{i} + 4\mathbf{j}$  respectively. Given that the system reduces to a couple, find  $\mathbf{F}$  and p.

Find the couple in the form  $a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$ .


12. A lamina in the shape of a rectangle ABCD with AB = a and BC = b is immersed in a homogeneous liquid with its surface vertical and the edge AB on the free surface of the liquid. Show that the centre of pressure of the lamina is at a depth  $\frac{2b}{3}$  from the free surface of the liquid.

A cubical tank has, on one of its vertical sides, a lid PQRS in the shape of a square of side of length c which is smoothly hinged along the horizontal top edge PQ. The tank is filled with a homogeneous liquid of density  $\rho$  to the level of PQ. Find the magnitude of the minimum force that must be applied at the mid-point of RS perpendicular to the plane of PQRS in order to keep the lid closed.


13. A particle P of mass m is projected vertically upwards under gravity with a initial velocity u. Assume that the resistance of air is  $gv^2$  per unit mass, where v is the velocity of P. Obtain the equation of motion of P and show that P comes to rest at a height  $\frac{1}{2g}\ln(1+u^2)$ .

Show that the velocity  $\omega$  with which the particle will return to the point of projection is given by  $\omega = \frac{u}{\sqrt{u^2 + 1}}$ .

**14.** (a)



A smooth sphere P moves with a speed u and makes a perfectly elastic collision with an identical sphere Q, which is at rest. Immediately before the collision, the velocity of P makes an angle  $\theta$  with the line of centres (see figure (1)). Find the speeds of P and Q after the collisions and show that the directions of motion of P and Q after the collisions are perpendicular.



In a billiard game, all the billiard balls are identical and of radius r. Assume that all the balls are smooth and the collisions are perfectly elastic.


Two red balls  $R_1$  and  $R_2$  are initially at rest with their centres at the points A and B respectively, where AB = d(>3r). A white ball W moves with a speed u towards  $R_1$  in a direction perpendicular to AB. After hitting  $R_1$ , W changes its direction and then collides with  $R_2$ . Suppose that at the moment of hitting  $R_2$ , the centre of W lies at a point C on AB (see figure (2)). Using the result in part(a), or otherwise,

- (i) state the directions in which  $R_2$  and W will move after the  $2^{nd}$  collision and
- (ii) find the speeds of  $R_2$  and W after the  $2^{nd}$  collision.
- 15. A frame consists of two uniform rods AB and BC, having lengths 4a and 2a respectively, which are rigidly joined together at B in such a way that  $A\hat{B}C = 60^{\circ}$ , (see the figure). The mass per unit length of both rods is m. Show that the moment of inertia of the frame about an axis through A perpendicular to the plane of the frame is  $48ma^3$ .

The frame can rotate freely about a fixed horizontal axis through A and perpendicular to the frame. Initially AB is vertical, with B above A, and the

frame is given an angular speed  $\sqrt{\frac{3g}{8a}}$ . The sense of the motion of the frame is such that, when AB is first horizontal, C is below AB. Show that angular

speed of the frame at the instant when AB is horizontal is  $\sqrt{\frac{(24+\sqrt{3})g}{24a}}$ 



- 16. A computer game consists of two levels, Level 1 and Level 2. A player starts at Level 1 and proceeds to Level 2, regardless of the outcome at Level 1. The probability of a win at Level 1 is 0.6. The probability of a win at Level 2 is 0.4 or 0.2, depending on a win at Level 1 or a loss at Level 1 respectively.
  - (i) Find the probability that a player
    - (a) wins at both levels,
    - (b) wins only one of the two levels.
  - (ii) Given that a player has won at Level 2 of a game, find the probability that he has also won at Level 1 of that game.
  - (iii) A player repeatedly plays this computer game until he wins at both levels of a game. Suppose that the outcomes of different games are independent. Find the expected number of games that a player has to play to win both levels of a game.
- 17. (a) A random variable X follows a Poisson distribution with the probability mass function  $P(X=x) = \frac{e^{-\lambda} \lambda^x}{x!} \text{ for } x = 0, 1, 2, ..., \text{ where } \lambda (>0) \text{ is a parameter.}$

Show that  $E(X) = \lambda$  and  $E(X^2) = \lambda^2 + \lambda$ .

The number of misrecognized letters per page by a character recognition system, follows a Poisson distribution with a mean of 0.5. Find the variance of the number of misrecognized letters in a page.

Given that this system has misrecognized at most two letters in a particular page, find the probability that there are no misrecognized letters in that page.

(b) Of a certain production, the copper content in a product follows a normal distribution with a mean of 59.9 grams and a standard deviation of 2.5 grams. Find the probability that the copper content of a randomly selected product from this production is more than 61 grams. Among the products with copper content less than 61 grams, find the percentage that will have copper content exceeding 60 grams.

සියලුම හිමිකම් ඇව්රිණි / $\psi$ ழுப் பதிப்புரிமையுடையது / $All\ Rights\ Reserved$ ]

ලි ලංකා විභාග දෙපාර්තමේන්තුව ලි ලංකා විභාග දෙපාර්තමේන්තුව කියලා දෙපාර්තමේන්තුව විභාග දෙපාර්තමේන්තුව ලි ලංකා විභාග දෙපාර්තමේන්තුව මි ලංකා විභාග දෙපාර්තමේන්තුව ලි ලික් දෙපාර්තමේන්තුව ලි ලංකා විභාග දෙපාර්තමේන්තුව ලි ලංකා විභාග දෙපාර්තමේන්තුව ලි ලික් දෙපාර්තමේන් ලි ලික් දෙපාර්තමේන්තුව ලි ලික් දෙපාර

අධායන පොදු සහතික පතු (උසස් පෙළ) විභාගය, 2024 கல்விப் பொதுத் தராதரப் பத்திர (உயர் தர)ப் பரீட்சை, 2024 General Certificate of Education (Adv. Level) Examination, 2024

උසස් ගණිතය உயர் கணிதம் Higher Mathematics



පැය තුනයි மூன்று மணித்தியாலம் Three hours

Use additional reading time to go through the question paper, select the questions you will answer and decide which of them you will prioritise.

Index Number	Ę				
Thuck I tulinger		200		 3° .	-

### **Instructions:**

- \* This question paper consists of two parts.

  Part A (Questions 1 10) and Part B (Questions 11 17).
- \* Part A:

  Answer all questions. Write your answers to each question in the space provided. You may use additional sheets if more space is needed.
- Answer five questions only. Write your answers on the sheets provided.
- \* At the end of the time allotted, tie the answer scripts of the two parts together so that Part A is on top of Part B and hand them over to the supervisor.
- \* You are permitted to remove only Part B of the question paper from the Examination Hall.

## For Examiners' Use only

(11) Higher Mathematics I					
Part	Question No.	Marks			
	1				
-	2				
	3	***************************************			
	4	· · · · · ·			
A	5				
	6				
	7				
	8				
	9				
	10				
	11				
	12				
ъ	13	,			
В	14				
	15	·			
	16				
	17				
	Total	· · · · · · · · · · · · · · · · · · ·			

	 Total
In Numbers	
In Words	

		Code Numbers
Marking Exami	ner	
Checked by:	1	
Checked by.	2	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
Supervised by:	4	

	Part A
. Fa	actorize: $a^3(b-c) + b^3(c-a) + c^3(a-b)$ .
4	
••	<u> </u>
••	
••	
• •	
••	
. • •	
•• نهري آه	
••	
• •	
a	nd 5. Show that $R$ is an equivalence relation on $\mathbb{Z}$ and write down the equivalence class of
•1	
•	
•	· · · · · · · · · · · · · · · · · · ·
:	
	A control of the cont
*	
•	
•	
•	
•	
•	
٠	
•	
•	

- 3 -

	lso find f	(1).								
• • •								• • • • • • • • • • • • • • • • • • • •		
		••••			• • • • • • • • • • • • • • • • • • • •		•••••		• • • • • • • • • • • • • •	
• • •	******	*********		•••••	· • • • • • • • • • • • • • • • • • • •		* * * * * * * * * * * * * * * * * * * *	• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •	
		•••••	* • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •	· • • • • • • • • • • • • • • • • • • •			• • • • • • • • • • • • • • • • • • • •		.,
• • •		*******			· • • • • • • • • • • • • • • • • • • •			.,		
									, , , , , , , , , , , , , , , , , , , ,	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
• • •	*********	*********	• • • • • • • • • • • •		•••••••		••••••			• • • • • • • • • • • • • • • • • • • •
•. • •.		******					• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •		,,
										Nata Kapar
	******						• • • • • • • • • • • • • • • • • • • •			• • • • • • • • • • • • • • • • • • • •
• • • •		•••••••••								
• • •		.,,,					• • • • • • • • • • • • • • • • • • • •		•	
	Section of the sectio		ing Single Single	era (jr. National)a				7 A.S.		
*****										
	•••••	• • • • • • • • • • • • • • • • • • • •				• • • • • • • • • • • • • • • • • • • •				
	5. + m <sup>2</sup>	a a a a a		2	3					
_				$\begin{vmatrix} a & a \end{vmatrix}$	1+a					
Let	t a, b, c	$\in \mathbb{R}$ . Sh	ow that	$ b  b^2$	$1+b^3$	= (1 + abc)	(a-b)(b	-c)(c-a)		1 - 4 - 4 - 4 - 4 - 4 - 4 - 4 - 4 - 4 -
			-	$c$ $c^2$	$1 + c^3$					13.
		-			ŀ					
••••	******	•••••••	• • • • • • • • • •	• • • • • • • • • •		• • • • • • • • • • • • • • • • • • • •				
••••										
									In the State of	
••••	**********				• • • • • • • • • • •	,				• • • • • • • • • • • • • • • • • • • •
		••••••••••••••••••••••••••••••••••••••					• • • • • • • • • • • • • •		*	
	**********		••••				• • • • • • • • • • • • • • • • • • • •			
			· · · · · · · · · · · · · · · · · · ·							
• • • •			×							
• • • •			v V							
			×							
••••		· · · · · · · · · · · · · · · · · · ·								
••••										

, , , , , , , , , , , , , , , , , , ,	AL/	2024/11/E-I -4-
	=	Show that the tangents drawn to the parabola $y^2 = 4\alpha x$ from the point $(-a, 0)$ are perpendicular
	3.	
l		
1	20 m 14 m 2 m	하게 하게 하다 사람들은 경에 집중을 가장하는 사람들이 들어가 되었다.
-		
-		
	ii (s	Let $a, b \in \mathbb{R}$ . Also, let $f: \mathbb{R} \to \mathbb{R}$ be the function given by $f(x) = \begin{cases} \frac{a(\sqrt{x+1}-2)}{x-3} & \text{if } x > 3, \\ x = 3 & \text{otherwise} \end{cases}$
	6.	Let $a, b \in \mathbb{R}$ . Also, let $f: \mathbb{R} \to \mathbb{R}$ be the function given by $f(x) = \begin{cases} x-3 \\ -3 \end{cases}$
		$e^{x-3} + bx  \text{if}  x \le 3.$
		It is given that $f$ continuous and the gradient of the tangent line drawn to the graph of $f$ when $x = is e^{-3} + 1$ . Find the values of $a$ and $b$ .
		· · · · · · · · · · · · · · · · · · ·
	-	
		, <u>,</u>

Now, let $h: \mathbb{R} \to \mathbb{R}$ be the function defined by $h(x) = \begin{cases} 2^x & \text{if } x \le 0, \\ 1 + \ln(x+1) & \text{if } x > 0. \end{cases}$ Is $h$ differentiable at $x = 0$ ? Justify your answer.	Find the gradient of the tangent lines drawn to the graphs of $f$ and $g$ when $x=0$ .						
Now, let $h: \mathbb{R} \to \mathbb{R}$ be the function defined by $h(x) = \begin{cases} 1 + \ln(x+1) & \text{if } x > 0. \end{cases}$ Is $h$ differentiable at $x = 0$ ? Justify your answer.	_	, v					
Is h differentiable at $x=0$ ? Justify your answer. $ \frac{1+\ln(x+1) \text{ if } x>0.}{1+\ln(x+1)} $ Is h differentiable at $x=0$ ? Justify your answer.	Nover let by D ->	D ha tha firmat	tion defined by	h(x) =			
is $h$ differentiable at $x=0$ ? Justify your answer.  Solve the differential equation $\frac{dy}{dx} = \frac{x-y}{x+y} + \frac{y}{x}$ .	Now, let n. K	N be the function	non defined by		n (c o		
Solve the differential equation $\frac{dy}{dx} = \frac{x-y}{x+y} + \frac{y}{x}$ .				$\int 1 + \ln(x +$	1) if $x > 0$ .		
Solve the differential equation $\frac{dy}{dx} = \frac{x-y}{x+y} + \frac{y}{x}$ .	s h differentiable	at $x = 0$ ? Justif	y your answer.			w	
Solve the differential equation $\frac{dy}{dx} = \frac{x-y}{x+y} + \frac{y}{x}$ .		•			e e		
Solve the differential equation $\frac{dy}{dx} = \frac{x-y}{x+y} + \frac{y}{x}$ .							
Solve the differential equation $\frac{dy}{dx} = \frac{x-y}{x+y} + \frac{y}{x}$ .							
Solve the differential equation $\frac{dy}{dx} = \frac{x-y}{x+y} + \frac{y}{x}$ .					* * * * * * * * * * * * * * * * * * * *		
Solve the differential equation $\frac{dy}{dx} = \frac{x-y}{x+y} + \frac{y}{x}$ .						,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	
Solve the differential equation $\frac{dy}{dx} = \frac{x-y}{x+y} + \frac{y}{x}$ .							
Solve the differential equation $\frac{dy}{dx} = \frac{x-y}{x+y} + \frac{y}{x}$ .							
Solve the differential equation $\frac{dy}{dx} = \frac{x-y}{x+y} + \frac{y}{x}$ .							
Solve the differential equation $\frac{dy}{dx} = \frac{x-y+y}{x+y+x}$ .							
Solve the differential equation $\frac{dy}{dx} = \frac{x-y+y}{x+y+x}$ .					.,		
Solve the differential equation $\frac{dy}{dx} = \frac{x-y+y}{x+y+x}$ .				* Land State Comment	الله الله الله الله الله الله الله الله		
Solve the differential equation $\frac{dy}{dx} = \frac{x - y}{x + y} + \frac{y}{x}$ .	<b>5</b> ************************************	နေရ ရေးရ ရေးရရှိနေရာသည်။ ချေရာခြားချစ်ချိန်းချိန်းကျားများကြားသည်။ ကြို့ကြို့	ৰ ভাৰত ভূমিক কুন স্থানিস্কৃতি কিবল কৰিব কিবল বিজ্ঞান				
Solve the differential equation $\frac{dy}{dx} = \frac{x-y}{x+y} + \frac{y}{x}$ .							
Solve the differential equation $\frac{dy}{dx} = \frac{x-y}{x+y} + \frac{y}{x}$ .							
				وبجيجه بجيد بجيديهم وبالج			
			3-1				
			. La Sala da La Sala Balbada Albada (n. 1879)		iki maj majako pilak ngilak naiki Ti		
	19-49-30-19-1 - v - #-0-6						
		. dy	x-y $y$				
	Solve the different	tial equation $\frac{1}{dx}$	$=\frac{1}{x+y}+\frac{1}{x}$ .	CONTRACTOR OF THE STATE OF THE	and the second of the second of the second		
		u.					
	y * - , ' - *	, <del>u</del>					
	•••••						
		· · · · · · · · · · · · · · · · · · ·					
		· · · · · · · · · · · · · · · · · · ·					
		· · · · · · · · · · · · · · · · · · ·					
		· · · · · · · · · · · · · · · · · · ·					
		· · · · · · · · · · · · · · · · · · ·					
		· · · · · · · · · · · · · · · · · · ·					
		· · · · · · · · · · · · · · · · · · ·					
		· · · · · · · · · · · · · · · · · · ·					
		· · · · · · · · · · · · · · · · · · ·					
		· · · · · · · · · · · · · · · · · · ·					
		· · · · · · · · · · · · · · · · · · ·					
		· · · · · · · · · · · · · · · · · · ·					
		· · · · · · · · · · · · · · · · · · ·					
		· · · · · · · · · · · · · · · · · · ·					
		· · · · · · · · · · · · · · · · · · ·					
		· · · · · · · · · · · · · · · · · · ·					
		· · · · · · · · · · · · · · · · · · ·					
· · · · · · · · · · · · · · · · · · ·		· · · · · · · · · · · · · · · · · · ·					
		· · · · · · · · · · · · · · · · · · ·					

ī.,		•			
Show that $\int \{x\}$	$f'(x) + x^2 f''(x) \} $	$dx = \int f(x)  dx$			
<b>j</b> ()		0	7		
v					
					·
			,		
		•••••			
					/ ·
**************************************					
			, , , , , , , , , , , , , , , , , , ,		
. <u> </u>					
gravita <del>na</del> se e e e e e e	er Artistanis (1985) Artistanis (1985)	and the second		ek ĝigla a grafija de a e e e e e e e	
خُو فَاوْدَ عِنْ عِنْ كُونَ فَالْمِثْ فِي فَالْمِثْ فِي فَالْمِثْ فِي فَا فَوْ عِنْ فِي فَالْمِثْ فِي فَالْمِ		د کار در	ાં મુખ્યત્વે કર્યા છે. મુખ્યત્વે મુખ્યત્વે	, ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ;	-
	on the first year.				
*******			• • • • • • • • • • • • • • • • • • • •		
					,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
	. 6				
		**************************************	ere e e e e e e e e e e e e e e e e e e		• • • • • • • • • • • • • • • • • • •
On the same d	liagram, sketch t	he curves wh	ose polar equat	ions are $r=1$	$+\sin\theta$ and $r=3$
On the same d	iagram, sketch t	he curves wh	ose polar equat	ions are $r = 1$ to curves with	$+\sin\theta$ and $r=3$ : $0 < \theta < \frac{\pi}{2}.$
On the same d	liagram, sketch tinates of the poi	he curves wh	ose polar equat	ions are $r = 1$ vo curves with	$+\sin\theta$ and $r=3$ : $0 < \theta < \frac{\pi}{2}.$
On the same d	iagram, sketch tinates of the poi	he curves wh	ose polar equat	ions are $r = 1$ to curves with	$+\sin\theta$ and $r=3$ : $0 < \theta < \frac{\pi}{2}.$
On the same d	iagram, sketch t inates of the poi	he curves wh	ose polar equat	ions are $r = 1$ to vo curves with	$+\sin\theta$ and $r=3$ : $0 < \theta < \frac{\pi}{2}.$
On the same d	iagram, sketch t inates of the poi	he curves what of intersec	ose polar equaterion of these to	ions are $r = 1 + 1$ wo curves with	$+\sin\theta$ and $r=3$ : $0 < \theta < \frac{\pi}{2}.$
On the same d	iagram, sketch t inates of the poi	he curves when tof intersect	ose polar equat	ions are $r = 1$ vo curves with	$+ \sin \theta \text{ and } r = 3 :$ $0 < \theta < \frac{\pi}{2}.$
On the same d	iagram, sketch t inates of the poi	he curves when tof intersec	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch t inates of the poi	he curves when tof intersection	ction of these tw	ions are $r = 1$ wo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch t inates of the poi	he curves when the of intersection	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch t inates of the poi	he curves when tof intersect	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch t	he curves wh	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch t	he curves wh	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch t	he curves wh	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch t	he curves wh	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch t	he curves wh	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch t	he curves wh	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch t	he curves wh	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch t	he curves wh	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch t	he curves wh	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch t	he curves when the of intersection of intersection intersection.	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch t	he curves when the of intersection of intersection intersection.	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch t	he curves wh	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch t	he curves wh	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch t	he curves whent of intersection	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch tinates of the poi	he curves when the of intersection of intersection intersection.	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch tinates of the poi	he curves when the of intersection of intersection intersection.	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same d	iagram, sketch t inates of the poi	he curves when the of intersection of intersection intersection.	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .
On the same difind the coordi	iagram, sketch t	he curves when the of intersection of intersection intersection.	ction of these tw	vo curves with	$0 < \theta < \frac{\pi}{2}$ .

கீகஓல் கிறிவர் ඇசிற்கி/முழுப் பதிப்புநிமையுடையது/ $All\ Rights\ Reserved$ ]

ලි ලංකා විහාත දෙපාර්තමේන්තුව ලී ලංකා විහාත දෙපාර්තමේ**සි වියාහ විභාග දෙපාර්තමේන්තුව** විහාත දෙපාර්තමේන්තුව ලී ලංකා විහාත දෙපාර්තමේන්තුව இலங்கைப் பழிட்சைத் திணைக்களம் இலங்கைப் பழிட்செத் திணைக்கதும் இருப்படைப் பழிட்சைத் திணைக்களம் இலங்கைப் பழிட்சைத் திணைக்களம் Department of Examinations, Sri Lanka Department of **இலங்கைப் Sri Lanka** ලී ලංකා විහාත දෙපාර්තමේන්තුව ලී ලංකා විභාග දෙපාර්තමේන්තුව ලී ලංකා විභාග දෙපාර්තමේන්තුව இலங்கைப் பழீட்சைத் திணைக்களம் இலங்கைப் பழீட்சைத் திணைக்களை

> අධායන පොදු සහතික පතු (උසස් පෙළ) විභාගය, 2024 සබාඛ්ධ பொதுத் தராதரப் பத்திர (உயர் தர)ப் பரீட்சை, 2024 General Certificate of Education (Adv. Level) Examination, 2024

උසස් ගණිතය I உயர் கணிதம் I Higher Mathematics I



# Part B

- \* Answer five questions only.
- 11. (a) Let A, B and C be subsets of a universal set S. Stating clearly the laws of Algebra of sets that you use, show that
  - (i)  $(B \cup C) \cap A' = (A' \cap B) \cup (C \cap A' \cap B')$
  - (ii)  $A \cap (B \cap C') = (A \cap B) \cap (A \cap C)'$ .
  - (b) Among a group of 400 tourists, 103 preferred to visit only Anuradhapura, 32 preferred to visit only Kandy and 71 preferred to visit only Jaffna. 19 preferred to visit all three cities. 41 do not prefer to visit any of these three cities. 235 do not prefer to visit Kandy. 200 do not prefer to visit Jaffna.

Find the number of tourists who preferred to visit only Kandy and Jaffna.

**12.** (a) Let  $a, b, c \in \mathbb{R}$  and a, b, c > 0.

Show that

(i) 
$$\frac{a+b}{2} \ge \sqrt{ab}$$
,

(ii) 
$$\frac{a^2+b^2}{c^2} + \frac{b^2+2c^2}{a^2} + \frac{2c^2+a^2}{b^2} \ge 2+4\sqrt{2}$$
.

(b) Find the value of  $\lambda$  such that,  $\begin{pmatrix} 3 & 2 \\ -8 & -5 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = \lambda \begin{pmatrix} u \\ v \end{pmatrix}$  for some non-zero  $u, v \in \mathbb{R}$ .

Hence or otherwise, find the equation of the line through the origin which gets mapped onto itself, under the transformation defined by  $\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} 3 & 2 \\ -8 & -5 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$ .

13. State and prove De Moivre's Theorem for a positive integral index.

Using De Moivre's theorem, show that  $\cos 5\theta = \cos^5 \theta - 10\cos^3 \theta \sin^2 \theta + 5\cos \theta \sin^4 \theta$  and  $\sin 5\theta = 5\cos^4 \theta \sin \theta - 10\cos^2 \theta \sin^3 \theta + \sin^5 \theta$  for  $\theta \in \mathbb{R}$ .

Hence, show that  $\tan 5\theta = \frac{5 \tan \theta - 10 \tan^3 \theta + \tan^5 \theta}{1 - 10 \tan^2 \theta + \tan^4 \theta}$  when  $\theta \neq \frac{n\pi}{5}$  for  $n \in \mathbb{Z}$ .

Prove that  $\tan \frac{\pi}{5}$  is a root of the equation  $x^4 - 10x^2 + 5 = 0$ .

Give the other roots of the equation  $x^4 - 10x^2 + 5 = 0$  in the form  $\tan \frac{k\pi}{5}$ , stating the values of k. **Deduce** that  $\tan^2 \frac{\pi}{5} + \tan^2 \frac{2\pi}{5} = 10$ .

14. (a) Let  $C_1$  and  $C_2$  be the curves given by  $y = |x^2 - 1|$  and  $y = x^2 - x|x|$  respectively. Sketch the region S, enclosed by the curves  $C_1$  and  $C_2$ .

Show that the area of the region S is  $\frac{2}{3\sqrt{3}}(1+\sqrt{3})$ .

Also, find the volume of the solid generated by rotating S about the x-axis through  $2\pi$  radians.

- (b) Solve the differential equation  $(x^2 + 1)\frac{dy}{dx} + xy = \sqrt{x^2 + 1}\sin x$  subject to the condition y = 1 when x = 0.
- **15.** (a) Let  $I_n = \int_0^{\infty} x^n \sqrt{a^2 x^2} \, dx$  for a > 0, and  $n = 0, 1, 2, 3, \dots$ Show that  $I_n = \frac{(n-1)}{(n+2)} a^2 I_{n-2}$  for  $n \ge 2$ .

Hence, evaluate  $\int_{0}^{2} x^{4} \sqrt{4 - x^{2}} dx.$ 

(b) Find the Maclaurin series of  $\sin x$  and  $\ln(1-x)$  in ascending powers of x upto and including the term in  $x^5$ .

**Hence**, obtain the Maclaurin series of  $\sin 2x \ln(1+2x)$  in ascending powers of x upto and including the term in  $x^3$ .

Using this, find an approximate value for  $\int_{0}^{4} \sin 2x \ln \left( \frac{1+2x}{1-2x} \right) dx.$ 

- **16.** (i) Prove that the line y = mx + c is a tangent to the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  if and only if  $c^2 = a^2m^2 + b^2$ .
  - (ii) Show that the locus of the point of intersection of the tangents to the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  which are at right angles, is given by  $x^2 + y^2 = a^2 + b^2$ .
  - (iii) A tangent to the ellipse  $x^2 + 4y^2 = 4$  meets the ellipse  $x^2 + 2y^2 = 6$  at the points P and Q. Show that the tangents at P and Q of the ellipse  $x^2 + 2y^2 = 6$  are at right angles.
- 17. (a) Let  $f(x) = \frac{1 \cos 2x}{3 + \cos 2x}$  for  $x \in \mathbb{R}$ .
  - (i) Show that  $0 \le f(x) \le 1$  for  $x \in \mathbb{R}$ .
  - (ii) Solve the equations f(x) = 0 and f(x) = 1, and sketch the graph of y = f(x) for  $0 \le x \le 2\pi$  indicating the turning points.
  - (b) The following table gives the values of  $f(x) = e^{-x^2}$ , correct to two decimal places, for values of x between 0 and 1:

*	0.00	0.25	0.50	0.75	1.00
$f(x) = e^{-x^2}$	1.00	0.94	0.78	0.57	0.37

Applying **Simpson's Rule**, find an approximate value for  $\int_{-x^2}^{1} dx$ .

An approximate value of  $\int_{0}^{1} \left(15 - e^{3-x^2}\right) dx$  is given to be 0.54. Find an approximate value for  $e^3$ .